



An Unusual Phenomenon of Gall Stones-Mirizzi Syndrome

Shaik AR*, Al-meer FM and Patel NK

Department of Family Medicine, Primary Health Care Corporation Doha, Qatar

Abstract

Most common cause of jaundice in primary care is biliary tree stones with a frequency of up to 33%. Mirizzi syndrome is a rare condition caused by the obstruction of the common bile duct or common hepatic duct by external compression from multiple impacted gallstones or a single large impacted gallstone in Hartman's pouch.

Introduction

Most common cause of jaundice in primary care is biliary tree stones with a frequency of up to 33% [1]. Mirizzi syndrome is a rare condition caused by the obstruction of the common bile duct or common hepatic duct by external compression from multiple impacted gallstones or a single large impacted gallstone in Hartman's pouch. Mirizzi syndrome without preoperative diagnosis could be as high as 17% [2]. Preoperative diagnosis of Mirizzi syndrome is difficult and can be made in only 8% to 62.5% of patients. It has been reported that 6% to 28% of patients with preoperative diagnosis of Mirizzi syndrome actually had gallbladder cancer [2]. Mirizzi syndrome is a condition difficult to diagnose and treat, representing a particular "challenge" for the biliary surgeon. The disease can mimic cancer of the gallbladder, causing considerable diagnostic difficulties [3].

Case Presentation

A 22-year-old well gentleman presented with 4 day's history of yellowish discoloration to eyes and skin. Additionally, he had abdominal pain on eating food and weight loss of 10 kg over past 6 months. Initial systemic physical examination was unremarkable except for icterus. He was referred to secondary care for further evaluation and management. Blood test confirmed deranged hyperbilirubinemia of 262 (Direct 217), Alkaline Phosphatase 386, ALT 267, AST 132.

Abdominal ultrasound showed thickened walls of the gallbladder with a calculus suggestive of cholecystitis. MRCP showed stones of the extrahepatic bile ducts causing intrahepatic bile duct obstruction, probably located within the lower end of the anomalously crossing cystic duct causing extrinsic compression of the common duct (Mirizzi Syndrome) a Laparoscopic Cholecystectomy + Trans cystic CBD exploration with Choledochoscope and extraction of stones by Dormia basket + Intraoperative Cholangiogram (IOC) was carried out. Findings were in keeping with Mirizzi syndrome with fistula communicating mid cystic to proximal CBD. IOC after the above procedure showed smooth passage of contrast to biliary tree. Blood tests done post operatively showed improvement in bilirubin (reducing to 125) and LFTs completely normalizing within a span of 20 days. He was discharged home on oral antibiotics and analgesia.

Discussion

Mirizzi syndrome is a rare complication caused by gallstone(s) located in the infundibulum of the gallbladder or cystic duct leading to adjacent biliary duct compression which results in partial or complete obstruction of the common hepatic duct and sometimes with the presence of cholecysto-choledochal fistula [4]. Incidence ranges from 1% per year in developed countries to up to 5.7% in developing countries. Most commonly presentations are with obstructive jaundice with right upper quadrant pain or dark urine. Endoscopic retrograde cholangiopancreatography is the best diagnostic modality which is an invasive procedure. Surgical cholecystectomy is considered as the treatment of choice [5].

Conclusion

A multispecialty approach must be taken for this condition. Accurate diagnosis with an accurate evaluation of the extent of this disease must be concluded by all involved. The surgeon is the main specialist for diagnosis and treatment but familiarity and a high degree of suspicion of this condition must be present by the initially treating primary caregiver or emergency physician. Delay in making

OPEN ACCESS

*Correspondence:

Shaik AR, Department of Family Medicine, Primary Health Care Corporation Doha, Al Mina Street, Second Ring Area Authority, Opp. Double Tree Hilton Hotel, Qatar, Tel: +974-50207492; E-mail: rashidshaik0376@gmail.com

Received Date: 09 Nov 2020

Accepted Date: 25 Nov 2020

Published Date: 04 Dec 2020

Citation:

Shaik AR, Al-meer FM, Patel NK. An Unusual Phenomenon of Gall Stones-Mirizzi Syndrome. *Ann Clin Case Rep.* 2020; 5: 1894.

ISSN: 2474-1655

Copyright © 2020 Shaik AR. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

an adequate diagnosis and treatment could lead to increased patient morbidity and mortality. Although definite cause of jaundice is difficult to be identified in the primary care, early recognition of jaundice and prompt referral to secondary care for further evaluation and management has been found to have favourable outcomes [6].

References

1. Taylor A, Stapley S, Hamilton W. Jaundice in primary care: A cohort study of adults aged >45 years using electronic medical records. *Fam Pract*. 2012;29(4):416-20.
2. Beltrán MA. Mirizzi syndrome: History, current knowledge and proposal of a simplified classification. *World J Gastroenterol*. 2012;18(34):4639-50.
3. Clemente G, Tringali A, De Rose AM, Panettieri E, Murazio M, Nuzzo G, et al. Mirizzi Syndrome: Diagnosis and management of a challenging biliary disease. *Can J Gastroenterol Hepatol*. 2018;2018:6962090.
4. Fachrull, Stephanie A, Mudjaddid E. Mirizzi's Syndrome. *Indonesian J Gastroenterol Hepatol Dig Endosc*. 2015;16(3).
5. Valderrama-Trevino AI, Granados-Romero JJ, Espejel-Deloiza M, Chernitzky-Camaño J, Barrera Mera B, Estrada-Mata AG, et al. Updates in Mirizzi syndrome. *Hepatobiliary Surg Nutr*. 2017;6(3):170-8.
6. Jones MW, Ferguson T. Mirizzi Syndrome. *Stat Pearls* [Internet]. 2020.