



Priapism in a Patient with COVID-19: A Case Report

Agulleiro Juan Cruz*, Lapenna Leandro, Koren Claudio, Labrador Jorge, Coronado Paul and Villalba Bachur Roberto

Urology Service, Complejo Hospital Churrucá-Visca, Argentina

Abstract

A pro-inflammatory response in patients with coronavirus disease provokes hypercoagulability and hyperviscosity, following an excessive liberation of acute phase reactants. Thromboembolic complications such as Deep Venous Thrombosis, Pulmonary Embolism, Stroke and Acute Coronary Syndrome were described from the beginning of the pandemic as possible manifestations of this new virus. Priapism is a medical urgency that demands quick assessment and timely treatment.

A case of priapism is presented in a patient undergoing hospitalization for COVID-19. He was treated with cavernosal blood aspiration, irrigated with 0.90% w/v saline solution and intracavernosal injection of the sympathomimetic agent (adrenaline), with favorable response.

Keywords: Priapism, COVID-19; Thrombosis

Introduction

Since 2019, the world has been under a global health crisis because of COVID-19, affecting over 160 million people to date [1]. As we already know, the respiratory system is the most compromised. Nonetheless, manifestations at gastrointestinal, neurological, dermatological, and cardiovascular levels have also been reported. Around 5% of the patients require Intensive Care Unit management, mainly due to Acute Respiratory Distress Syndrome [2].

Within this group of patients, 31% suffer thromboembolic complications [3], with potentially fatal consequences such as: Deep Venous Thrombosis, Pulmonary Thromboembolism, Stroke and Acute Coronary Syndromes.

This case report presents a patient on his second day after admission because of COVID-19 infection, with priapism as a urogenital pathology. This finding aims at adding diagnostic and therapeutic evidence to the few cases already reported [4-6].

Case Presentation

A 62-year-old man with history of Arterial Hypertension, Mellitus Diabetes type 2, Grade 2 Obesity, and no urological history was admitted to the emergency room with febrile syndrome, headache, non-productive cough, anosmia, dysgeusia, myalgia, and generalized arthralgia. He had developed an unfavorable response, including respiratory failure. Supplemental oxygen was supplied at 10 L/min with a reservoir mask, and Intensive Care Unit management was required. COVID-19 diagnosis was confirmed by real-time reverse transcriptase-polymerase chain reaction on a nasopharyngeal swab specimen.

He then experienced hemodynamic failure, with supplemental O₂ at 15 L/min, unstable respiratory mechanics, tachypnea and respiratory failure; therefore, orotracheal intubation and mechanical respiratory assistance were established. (FIO₂ 60% and PaO₂/FiO₂ ratio of 85) (Figure 1).

Chest-Computed Tomography (CT) and CT angiography were performed at the initial admission and showed diffuse and bilateral involvement of the pulmonary parenchyma, with some areas indicating ground-glass opacity. No signs of proximal pulmonary embolism were identified.

After 24 h of admission into the ICU, the patient presented a 5-h episode of priapism. A consequent increase in D-dimer was observed, among other acute phase reactants, which expressed its maximum value on the fourth day with a level of 4,885, requiring anticoagulation, enoxaparin twice daily (BID) at a dose of 40 mg [7]. The patient was sedated and unable to answer questions on his medical record and pain levels. He had no previous history of priapism, he was not receiving any treatment for erectile sexual dysfunction, there was no trauma associated, and neither medication

OPEN ACCESS

*Correspondence:

Agulleiro Juan Cruz, Urology Service,
Complejo Hospital Churrucá-Visca,
Guardia vieja 4129, CABA, Argentina,
Tel: +5491168842019;
E-mail: juancruzagulleiro@hotmail.com

Received Date: 01 Nov 2021

Accepted Date: 08 Dec 2021

Published Date: 27 Dec 2021

Citation:

Cruz AJ, Leandro L, Claudio K, Jorge L, Paul C, Roberto VB. Priapism in a Patient with COVID-19: A Case Report. *Ann Clin Case Rep.* 2021; 6: 2070.

ISSN: 2474-1655

Copyright © 2021 Agulleiro Juan Cruz. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Clinical laboratory results		
	Patient	VN
Hematocrit, %	36	39-53
Leukocytes 10x3/mm3	22	<10
Hemoglobin, g/dL	13	13-18
Platelets, G/L	203	150-400
Urea, mmol/L	2	3.0-9.2
Creatinine, µmol/L	0.63	62-106
Glucose, mg /dL	106	70-140
Total Billirubin, µmol/L	11	< 21
D-Dimer ng/ml	885.	>255
Procalcitonin	0.22.	>0.005
Creatine phosphokinase	440.	>190
Lactate dehydrogenase	310	100-250
Prothrombin %	100	70-130

Fig 1. Clinical laboratory results at the time of priapism. VN = Normal Value

Figure 1: Clinical laboratory results at the time of priapism (VN: Normal Value).

Blood gas analysis (FiO2 60%)		
	Paciente	VN
PH	7.36	7.35-7.45
PaCO2mmHG	52	35-45
PaO2mmHG	66	85-100
HCO3- mEq/L	30	22-29
O2%	89%	95-98%

Cavernosal blood gas analysis	
PH	7.2
PaCO2 mmHg	80
PaO2 mmHg	60
CO2 total mmol/L	36
Lactate, mmol/L	8

Fig 2.A) Blood gas analysis at the time of priapism. Opposite mixed disorder is evidenced. B) Cavernosal blood gas analysis. CO2, carbon dioxide; O2, oxygen; PaCO2, partial pressure of CO2 in arterial blood; PaO2, partial pressure of O2 in arterial blood O2%, Oxygen saturation in blood.

Figure 2: Blood gas analysis at the time priapism. Opposite mixed disorder is evidenced. B) Cavernosal blood gas analysis. CO2: Carbon Dioxide; O2: Oxygen; PaCO2: Partial Pressue of CO2 in Arterial blood; PaO2: Partial Pressure of O2 in Arterial Blood; O2%: Oxygen Saturation in Blood

nor current disease could be identified as an etiologic agent of priapism [8].

Physical examination revealed induration of the corpora cavernosa with a flaccid glans. Given the persistence of the erection, the lack of response to local ice, and the inability to perform exercises to reduce priapism, it was decided to continue with the stepped care.

Two Abbocath 18 catheters, one in each corpora cavernosa, were

placed at 3 and 9 o'clock and dark blood with clots appeared. Thus, 4 ml of this dark blood were drawn and cavernosal blood gases analysis showed a venous sample with a pattern of metabolic acidosis (Figure 2). This finding served for a clinical diagnostic suspicion of Ischemic, Low flow or Veno-Occlusive priapism.

The patient was irrigated two times with 0.90% w/v saline solution at 20-min intervals, with no satisfactory response. The next therapeutic step was an injection in each corpora cavernosa with 1 ml of adrenaline (dilution of 1/100 ml w/v saline solution) at 20-min intervals, with no satisfactory response.

The therapeutic process was repeated. After 20 min, a decrease in the erection was confirmed. A compressive bandage and local ice were placed to avoid bruising. The patient did not repeat episodes of priapism in the following days.

Discussion

Priapism is defined as a partial or complete erection, with or without sexual stimulation that persists for more than 4 h, which can be painful. According to its pathophysiology, it can be classified into 3 types:

- Ischemic, low-flow or veno-occlusive
- Non-ischemic, high-flow, or arterial.
- "Stuttering", recurring or intermittent.

In this case, we will focus on Ischemic priapism. Ischemic priapism presents as rigidity of the corpora cavernosa associated with a flaccid state of the glans penis. It represents >95% of the cases. It is a urological emergency that requires quick assessment and timely treatment. If the erection persists over time, the smooth muscle cells suffer necrosis, producing fibrosis of the corpora cavernosa, leading the patient to permanent erectile sexual dysfunction. The most relevant data from the clinical history of a patient with priapism include: Duration of erection, presence and severity of pain, previous episodes of priapism and treatment provided, whether the patient is undergoing any treatment for erectile sexual dysfunction, medication or related drugs, hematological diseases (sickle cell disease, aplastic anemia, agranulocytosis, thrombocytopenia, and paraneoplastic syndromes, among others).

Focusing on the physiopathology of this condition, we find different metabolic alterations depending on the time of evolution. The smooth muscle cells of the corpora cavernosa experience hypoxia, hypercapnia, glyopenia and metabolic acidosis [9-10].

A histological analysis with serial biopsies of the muscle cells of the corpora cavernosa during an episode of priapism shows predominantly interstitial edema at 12 o'clock. After 24 h, destruction of the sinusoidal endothelium is mainly observed, exposing the basement membrane and revealing adhesion of platelets. After 48 h, irreversible changes appear in the cells of the corpora cavernosa with necrosis and scar tissue [11].

In the case of the patient presented, the erection was maintained for over 5 h, the presence of pain could not be verified (because of the patient's condition) and there was no etiological factor for priapism in the clinical history.

The type of erection, already described above, the color of the blood and the arterial gas blood test results strongly bring us closer to the clinical suspicion of ischemic priapism. The only finding that

could be considered discordant was the presence of a partial pressure of $O_2 > 30$ mmHg in the gases of the corpora cavernosa. There could be a correlation between this finding and the fact that the patient was intubated with 60% FIO₂ at the time.

It is already known that COVID-19 causes a pro-inflammatory state in patients, which also favors a pro-thrombotic state; and this is evidenced by the presence of hyperviscosity, hypercoagulability, favored by leukocytosis and endothelial dysfunction generated mainly by the release of interleukins and nitric oxide. This leads to a prothrombotic state, plus endothelial dysfunction. Some of these factors may trigger priapism.

Conclusion

The existence of the pandemic pushes health care workers of all specialties to the limit, since it demands maximal physical and mental effort. The multidisciplinary approach to this medical emergency allows us to carry out a fast and effective treatment, which, if not established in a timely manner, may lead to permanent sequela.

Limitations

Due to the limited number of published cases, evidence to confirm that COVID-19 virus is an etiological agent of priapism is not conclusive.

Acknowledgment

Urology Service from the Medical Complex of the federal police Churruca-Visca Hospital. Intensive Care Unit service from the Medical Complex of the federal police Churruca-Visca Hospital.

References

1. Johns Hopkins University (Baltimore, USA), local authorities last update of figures May 24, 2021;8:39 GMT.
2. Liang W, Liang H, Ou L, Chen B, Chen A, Li C, et al. Development and validation of a clinical risk score to predict the occurrence of critical illness in hospitalized patients with COVID-19. *JAMA Intern Med.* 2020;180(8):1081-9.
3. Klok FA, Kruip M, van der Meer NJM. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res.* 2020;191:145-7.
4. Lamamri M, Chebbi A, Mamane J, Abbad S, Munuzzolini M, Sarfati F, et al. Priapism in a patient with Coronavirus Disease 2019 (COVID-19). *Am J Emerg Med.* 2021;39:251.e5-251.e7.
5. Addar A, Al Fraidi O, Nazer A, Althonayan N, Ghazwani Y. Priapism for 10 days in a patient with SARS- CoV-2 pneumonia: A case report. *J Surg Case Rep.* 2021;2021(4):rjab020.
6. Silverman ML, Van Der Veer SJ, Donnelly TJ. Priapism in COVID-19: A thromboembolic complication. *Am J Emerg Med.* 2021;45:686.e5-686.e6.
7. COVID-19 and VTE-Anticoagulation. *Hematology.* 2020.
8. EUA guidelines on Sexual and Reproductive Health. 2020.
9. El-Bahnasawy MS, Dawood A, Farouk A. Low-flow priapism: Risk factors for erectile dysfunction. *BJU Int.* 2002;89(3):285-90.
10. Muneer A, Cellek S, Dogan A, Kell PD, Ralph DJ, Minhas S. Investigation of cavernosal smooth muscle dysfunction in low flow priapism using an in vitro model. *Int J Impot Res.* 2005;17(1):10-8.
11. Vreugdenhil S, Jorge PJF, van Driel MF, Nijsten MW. Ischemic priapism as a model of exhausted metabolism. *Physiol Rep.* 2019;7(6):e13999.