



## Shrimp Allergy: More and More Interesting

Riccardo Asero\*

Ambulatorio di Allergologia, Clinica San Carlo, Italy

### Commentary

In recent years food allergy has elicited growing attention on the media and in the general public. Crustaceans represent one of the most prevalent causes of food allergy worldwide, and in Italian adults they are the second cause of primary food allergy and food-induced anaphylaxis, after plant-derived foods [1,2]. Not many years ago it was generally believed that the only shrimp allergen was tropomyosin, a phylogenetically conserved muscle protein showing a highly homologous sequence in all invertebrates (worms, crustaceans, mollusks, snails, and insects) [3]. However, the complexity of the allergenic profile of shrimp has been increasingly recognized over the last decade, and it now includes no less than 15 allergens some of which are species-specific while others are homologous, and hence cross-reacting, to allergens in house dust mites, other shellfish, and parasites like *Anisakis simplex*. Many of these allergens have not been characterized and sequenced yet. In a recent study, less than 50% of Italian shrimp-allergic patients were found to react to tropomyosin, and most of them were found to produce IgE specific for a variable number of different proteins [4]. Thus, the story of shrimp allergy is becoming more and more challenging but also increasingly stimulating. The physical and chemical features of shrimp allergens may differ, and this can translate into different clinical expressions of allergy. For instance, most allergens are heat- or pepsin-resistant but others are destroyed by heat or by pepsin digestion. As a consequence, some allergic patients may experience only local symptoms (e.g., oral allergy syndrome, asthma following inhalation of aerosols during cooking or industrial processing, or contact urticaria after handling shrimps), or react only following the ingestion of raw shrimp, while others will show systemic reactions following cooked shrimps.

Altogether, there is a lot of material to work on during the next years in order to better characterize this type of food allergy.

### References

1. Asero R, Antonicelli L, Arena A, Bommarito L, Caruso B, Crivellaro M et al. EpidemAAITO: Features of food allergy in Italian adults attending allergy clinics: a multicenter study. *Clin Exp Allergy*. 2009; 39: 547-55.
2. Asero R, Antonicelli L, Arena A, Bommarito L, Caruso B, Colombo G, et al. Causes of food-induced anaphylaxis in Italian adults: a multi-centre study. *Int Arch Allergy Immunol*. 2009; 150: 271-227.
3. Leung PS, Chu KH, Chow WK, Ansari A, Bandea CI, Kwan HS, et al. Cloning, expression, and primary structure of *Metapenaeus ensis* tropomyosin, the major heat-stable shrimp allergen. *J Allergy Clin Immunol*. 1994; 94: 882-890.
4. Asero R, Mistrello G, Amato S, Ariano R, Colombo G, Conte ME, et al. Shrimp allergy in Italian adults: a multicenter study showing a high prevalence of sensitivity to novel high molecular weight allergens. *Int Arch Allergy Immunol*. 2012; 157: 3-10.

### OPEN ACCESS

#### \*Correspondence:

Riccardo Asero, Ambulatorio di  
Allergologia, Clinica San Carlo, Via  
Ospedale, 21, 20037 Paderno Dugnano  
MI, Italy,

E-mail: r.asero@libero.it

Received Date: 27 Aug 2017

Accepted Date: 24 Oct 2017

Published Date: 26 Oct 2017

#### Citation:

Asero R. Shrimp Allergy: More and  
More Interesting. *Ann Clin Case Rep*.  
2017; 2: 1457.

ISSN: 2474-1655

Copyright © 2017 Riccardo Asero. This  
is an open access article distributed  
under the Creative Commons Attribution  
License, which permits unrestricted  
use, distribution, and reproduction in  
any medium, provided the original work  
is properly cited.