



Large Vessel Vasculitis - Evaluation with 18F-FDG PET-CT

Sousa V¹, Loewenthal C² and Vieira MR^{1*}

¹Department of Nuclear Medicine, Hospital da Luz, Portugal

²Department of Clinical Research, UCL Eastman Dental Institute, UK

³Department of Microbial Diseases, UCL Eastman Dental Institute, UK

Clinical Image

72 yr old female, past history of headache, hospitalized with abdominal pain, nausea, vomiting and diarrhea. No other signs or symptoms. Blood tests showed anaemia (Hg 9.7 g/dL), high CRP (9.7 mg/dL) and ESR (111 mm/h). Abdominal CT showed ascitis. Stool cultures, *Clostridium* toxin, colonoscopy, enteroscopy and biopsies were all negative. She was released 6 days later having improved clinically. Reevaluation one month later showed anaemia, high CRP and ESR. 18F-FDG PET-CT documented diffuse increased metabolism (SUVmax 15) in thoracic and abdominal aorta, iliac, femoral, subclavian, carotid and vertebral arteries, suggesting large vessel vasculitis (Figure 1). Therapy with prednisolone and azathioprine was maintained for four years, with clinical and analytic improvement. Three months after therapy withdrawal the 18FFDG PET-CT remained normal (Figure 2).




Figure 1: 18F-FDG PET-CT documented diffuse increased metabolism (SUVmax 15) in thoracic and abdominal aorta, iliac, femoral, subclavian, carotid and vertebral arteries, suggesting large vessel vasculitis.

OPEN ACCESS

*Correspondence:

Vieira MR, Department of Nuclear Medicine, Hospital da Luz, Av. Lusíada n. 100, 1500-650 Lisbon, Portugal, E-mail: rvieira@hospitaldaluz.pt

Received Date: 04 Oct 2016

Accepted Date: 11 Oct 2016

Published Date: 24 Oct 2016

Citation:

Sousa V, Loewenthal C, Vieira MR. Large Vessel Vasculitis - Evaluation with 18F-FDG PET-CT. *Ann Clin Case Rep.* 2016; 1: 1164.

Copyright © 2016 Vieira MR. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Figure 2: Three months after therapy withdrawal the 18FFDG PET-CT remained normal.