



Scrofuloderma Patient Who Treated Long -Time as Hydradenitis Suppurativa

Ayhan Bekmez*

Department of Biotechnology, Ankara University, Turkey

Keywords

Scrofuloderma; Hydradenitis suppurativa; Differential diagnosis

Clinical Image

27 -year-old male patient admitted to our out-patient clinic with depressed scarring granulomatous pustular lesion on the face, beard-region (Figure 1). He described hydradenitis suppurativa lesions axillary and inguinal folds and treated with antibiotics for 2 years but moderate improvement. The disease going on with new pustules (Figure 2). We suspected of scrofuloderma and sent our patient to tuberculosis fight center. PPD was 15 mm and lung graph was normal. The patient treated combined therapy for skin tuberculosis for 6 months. Scrofuloderma must be thought in the differential diagnosis of hydradenitis suppurativa [1-3].



Figure 1: Depressed scarring granulomatous and pustular lesions on the beard-region.

OPEN ACCESS

*Correspondence:

Ayhan Bekmez, Department of
Biotechnology, Ankara University,
Ankara, Turkey,

E-mail: ayhanbekmez@hotmail.com.tr

Received Date: 18 Feb 2019

Accepted Date: 26 Mar 2019

Published Date: 01 Apr 2019

Citation:

Bekmez A. Scrofuloderma Patient Who
Treated Long -Time as Hydradenitis
Suppurativa. Ann Clin Case Rep. 2019;
4: 1631.

ISSN: 2474-1655

Copyright © 2019 Ayhan Bekmez. This
is an open access article distributed
under the Creative Commons Attribution
License, which permits unrestricted
use, distribution, and reproduction in
any medium, provided the original work
is properly cited.



Figure 2: Axillary fold granulomatous and pustular lesions.

References

1. Franko-Parades C, Marcos LA, Henao-Martinez AF, Rodríguez-Morales AJ, Villamil-Gómez WE, Gotuzzo E, et al. Cutaneous mycobacterial infections. Clin Microbiol Rev. 2018;32(1).
2. Tadek H. Scrofuloderma with disseminated tuberculosis in an Ethiopian child : a case report. J Med case rep. 2018;12(1):371.
3. Mello RB, Vale ESCP, Baeta IGR. Scrofuloderma: a diagnostic challenge. An Bras Dermatol. 2019;94(1):102-4.