



Sigmoid Intussusception Secondary to Lipoma: Comprehensive Case Report and Review

Mohamed Ben Lahsen^{1,2*}, Hamza Sekkat^{1,2}, Youness Bakali^{1,2}, Farid Sabbah^{1,2}, Mohammed Raiss^{1,2}, Abdelmalek Hrra^{1,2} and Mouna EL Alaoui Mhamdi^{1,2}

¹Digestive Surgical Department C, Ibn Sina University Hospital, Rabat, Morocco

²Faculty of Medicine and Pharmacy, Mohammed V University in Rabat, Morocco

Abstract

Introduction: Intussusception refers to the invagination of one segment of the intestine into another, often leading to obstruction and necessitating urgent management, particularly in adult presentations.

It is rare in adults and most often secondary to an underlying cause, but the exact mechanism remains undefined.

It is classified based on location and etiology. Malignant pathology is the most common, particularly for colonic intestinal invagination.

Lipoma is a rare cause of intestinal invagination

Case Report: In this article, we report the rare case of a patient who presented with sigmoid intestinal invagination due to a benign adipose tumor, treated by surgical resection.

Discussion: Intestinal invagination in adults is different from that in children, which is most often idiopathic.

Malignant tumor pathology represents two-thirds of colonic intestinal invaginations. Colonic benign adipose tumor is a rare cause of intestinal invagination; the diagnosis is established through CT scans and colonoscopy, but it is often difficult to differentiate it from cancer.

This is why the case we present is interesting due to its location in the sigmoid colon, which is not common, and the diagnostic uncertainty poses a challenge for therapeutic management.

Conclusion: The management of colonic intestinal invagination in adults is not universally accepted; in cases of diagnostic uncertainty regarding the histological nature of the causal pathology, en bloc surgical resection without prior disinvagination seems to be the most appropriate approach given the high risk of malignancy.

Keywords: Adult intussusception, Sigmoidal benign adipose tumor, Management, Surgery

Introduction

Intussusception refers to the invagination of one segment of the intestine into another, often leading to obstruction and necessitating urgent management, particularly in adult presentations.

The exact mechanism which precipitates intestinal invagination intestine is still unknown, but it is generally believed that any lesion in the bowel wall or irritant normal within the bowel lumen which alters the peristaltic pattern is capable of starting an invagination [1,2].

A lead point is common and can be found in 70–90% of adult intestinal invaginations, contrary to pediatric intestinal invagination, which is idiopathic in 90% of cases [3].

Intussusception is classified based on location, as well as malignant, benign, or idiopathic etiology.

For invaginations secondary to a benign etiology, benign adipose tumor ranks third after hyperplastic and adenomatous polyps [3].

Lipoma is the most common nonepithelial benign tumor that occurs in the colon; it makes up 65% of the benign adipose tumors of the gastrointestinal tract. The incidence of benign adipose

OPEN ACCESS

*Correspondence:

Mohamed Ben Lahsen, Digestive Surgical Department C, Ibn Sina University Hospital, Rabat, Morocco, Tel: +212 650680618

Received Date: 14 Oct 2025

Accepted Date: 27 Oct 2025

Published Date: 28 Oct 2025

Citation:

Lahsen MB, Sekkat H, Bakali Y, Sabbah F, Raiss M, Hrra A, et al. Sigmoid Intussusception Secondary to Lipoma: Comprehensive Case Report and Review. *Ann Clin Case Rep.* 2025; 10: 2788.

ISSN: 2474-1655.

Copyright © 2025 Mohamed Ben Lahsen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1 (A, B): Axial CT scan images showing the sigmoid-to-sigmoid intestinal invagination with diffuse wall thickening.

tumors of the colon in routine autopsy of all cases has been reported to vary between 0,035 to 4,4% [4].

Here, we report the case of a 70-year-old female patient who presented with sigmoid intestinal invagination due to a benign adipose tumor, treated by surgical resection.

Case Presentation

This is a 70-year-old female patient with no significant medical history, who presented with recurrent subocclusive syndromes that resolved spontaneously over the past 3 months, associated with episodes of mild rectal bleeding. There were no other associated symptoms, notably no abdominal pain or vomiting. Clinical examination was unremarkable except for slight tenderness in the left iliac fossa. All of this occurred in the context of preserved general condition.

A colonoscopy was performed and revealed a large pedunculated polypoid formation with endoluminal development located 20 cm from the anal margin, obstructing the lumen. The pathological study showed no specific or tumoral lesion within the limits of the communicated sample.

The patient also underwent an abdominal-pelvic CT scan, which revealed a sigmoid-to-sigmoid and meso-sigmoid intestinal invagination, creating a target sign appearance associated with diffuse wall thickening, micronodular infiltration of the perilesional fat, and infiltration of the mesorectal fat (Figure 1).

The biological assessment was normal, with no anemia or biological inflammatory syndrome, and the tumor markers were negative.

The patient was presented at a multidisciplinary consultation meeting, and given the persistence of symptoms and the potential malignancy of the mass causing the intestinal invagination, the decision to operate on the patient was made.

The patient was approached laparoscopically, and intraoperative exploration revealed a large invaginated mass in the sigmoid loop measuring approximately 8 cm, associated with colonic and ileal dilation upstream, making dissection under laparoscopy difficult, hence the decision to convert. There were no other lesions, notably no hepatic lesions or peritoneal carcinomatosis nodules.

Despite the preoperative pathological examination which did not show signs of malignancy, and given the suspicious macroscopic appearance of the tumor (Figure 2), we decided to perform a carcinological resection of the tumor. Thus, a low segmental left colectomy was performed, respecting safety margins of 5 cm on either side of the invaginated mass without prior disinvagination,



Figure 2: Surgical specimen of low segmental left colectomy removing the intestinal invagination.

with lymph node dissection at the base of the sigmoid trunk; and the continuity was restored by a mechanical colorectal anastomosis using a circular stapler.

The postoperative course was uncomplicated, with good clinical and biological progress. Bowel function resumed on day 3, and the patient was able to start eating again on day 4. She was discharged on day 5.

The pathological examination of the surgical specimen revealed a remodeled colonic benign adipose tumor.

With a follow-up of 4 months, the patient is doing well and no longer reports subocclusions or rectal bleeding. A follow-up colonoscopy is scheduled 6 months postoperatively to explore the rest of the colon.

Discussion

Adult intestinal invagination is entirely different from the pediatric form in its clinical, pathological, and therapeutic aspects. Pediatric intestinal invagination is a common disease, generally idiopathic without an identified lead point. The clinical presentation is often acute and easily recognized by the classic triad of abdominal pain, rectal bleeding, and a palpable abdominal mass. Most cases can be successfully treated by hydrostatic or pneumatic reduction without surgical intervention [5].

On the other hand, adult intestinal invagination is a rare disease, accounting for only 1% of intestinal obstructions and nearly 5% of all intestinal invagination cases [6-8].

Depending on the location, intestinal invaginations are classified as ileo-ileal, ileo-colic, and colo-colic [9]. Colonic intestinal

invagination is rarer and accounts for 19.9% of all intestinal invaginations [10].

According to the etiology of adult intestinal invagination, the rates of malignant tumors, benign tumors, and idiopathic causes are 32.9%, 37.4%, and 15.1% respectively [10].

Two thirds of adult colo-colonic intestinal invaginations are secondary to a primary colonic cancer. The remaining third are secondary to polyps, adenomas, endometriosis, previous anastomosis and benign adipose tumors [11].

Lipoma is the third most common benign tumor after hyperplastic and adenomatous polyps and the most nonepithelial benign tumor that occurs in the colon; the majority (90%) are submucosal in location, the remaining ones are subserosal in location and related to the appendices epiploicae [12].

benign adipose tumor of the colon is predominantly localized in the ascending colon (61% of cases), followed by the descending colon (20.1%), the transverse colon (15.5%) and the rectum (3.4%) [13].

They are more common in women with a peak incidence between 50 and 60 years of age [14].

Symptoms, when present, seem to be related to the size of the benign adipose tumor. Lipomas less than 2 cm in diameter rarely produce symptoms, whereas benign adipose tumors greater than 2 cm in diameter are more likely to be symptomatic. Fifty percent of the patients with benign adipose tumors 3 to 4 cm in diameter have symptoms, and if the benign adipose tumor is greater than 4 cm in diameter, 75% of the patients have symptoms [12].

The most common symptoms are abdominal pain followed by change in bowel habits, bleeding, signs of obstruction, signs of intestinal invagination, or, rarely, a palpable mass [12].

The diagnosis can be made by imaging methods, particularly CT scans, which show a fatty density mass, but its performance is reduced for benign adipose tumors smaller than 2 cm. Colonoscopy allows direct visualization of the tumor, and biopsies can confirm the histological nature of the benign adipose tumor [14].

However, the diagnosis of benign adipose tumor usually is not made preoperatively. These lesions are often confused with carcinoma and adenomas because of the age range of patients and the similarity in the roentgenologic appearance [12].

Until now, there is no universally accepted approach to the optimal treatment of adult intestinal invaginations. Given the large proportion of underlying pathological lesions, surgical management continues to be the mainstay treatment modality for most patients, and en bloc resection without reduction is recommended for colonic intestinal invaginations considering the high rate of malignancy [15-17].

Conclusion

Adult intestinal invagination is relatively rare compared to that in children and is almost always secondary to an underlying cause.

The combination of colonoscopy and CT scan is an interesting method for positive and etiological diagnosis.

However, the high frequency of malignant etiologies of intestinal invagination, particularly colonic, and the difficulty in establishing

a differential diagnosis with benign causes pose a real challenge in managing this pathology.

In the absence of a certain diagnosis, surgical treatment by en bloc resection without prior disinvagination seems to be the best option given the high risk of malignancy.

Author Contribution

FS, AH, MR designed the paper. BM collected the data. HS and BM wrote the first draft of the manuscript. HS participated in the article design and critically reviewed the manuscript. BY, MAM Was responsible for the microscopic examination and the histopathological analysis of the specimen, AH critically reviewed the manuscript. All authors approved the final version of the manuscript.

References

- Zubaidi A, Al-Saif F, Silverman R. Adult Intussusception: A Retrospective Review. *Dis Colon Rectum*. 2006;49:1546–51.
- D Weilbaecher, J A Bolin, D Hearn, W Ogden 2nd. Intussusception in adults. Review of 160 cases. *Am J Surg*. 1971;121(5):531-5.
- Kansoun A, Al Mohtar I, Bahmad M, Houcheimi F, Maanieh N, Hazim M, et al. Colo-colic intussusception secondary to colon lipoma: A case report. *Int J Surg Case Rep*. 2021;81:105695.
- Gordon RS, O'Dell KB, Namon AJ, Becker LB. Intussusception in the adult--a rare disease. *J Emerg Med*. 1991;9(5):337-42.
- Tarchouli M, Ali AA. Adult Intussusception: An Uncommon Condition and Challenging Management. *Visc Med*. 2021;37:120–7.
- Azar T, Berger DL. Adult intussusception. *Ann Surg*. 1997;226(2):134–8.
- Gomes A, Sousa M, Pignatelli N, Nunes V. Adult intussusception: a single-center 10-year experience. *Eur Surg*. 2013;45(5):239–44.
- Traoré D, Sissoko F, Ongoïba N, Traoré I, Traoré AK, Koumaré AK. Adult intussusception: diagnostic pitfalls, morbidity and mortality in a developing country. *J Visc Surg*. 2012;149(3):e211–4.
- Nagorney DM, Sarr MG, McIlrath DC. Surgical management of intussusception in the adult. *Ann Surg*. 1981;193(2):230–6.
- Hong KD, Kim J, Ji W, Wexner SD. Adult intussusception: a systematic review and meta analysis. *Tech Coloproctol*. 2019;23(4):315–24.
- Takeuchi K, Tsuzuki Y, Ando T, Sekihara M, Hara T, Kori T, et al. The Diagnosis and Treatment of Adult Intussusception: *J Clin Gastroenterol*. 2003;36(1):1821.
- Gordon RT, Beal JM. Lipoma of the Colon. *Arch Surg*. 1978;113(7):897-9.
- Saba RB, Sadeghi A, Rad N, Safari MT, Barzegar F. Colonic intussusception in descending colon: An unusual presentation of colon lipoma. *Gastroenterol Hepatol Bed Bench*. 2016;9:S93–S96.
- M'rabet S, Jarrar MS, Akkari I, Abdelkader AB, Sriha B, Hamila F, et al. Colonic intussusception caused by a sigmoidal lipoma: A Case Report. *Int J Surg Case Rep*. 2018;50:1-4.
- Goh BK, Quah HM, Chow PK, Tan KY, Tay KH, Eu KW, et al. Predictive factors of malignancy in adults with intussusception. *World J Surg*. 2006;30(7):1300–4.
- Kim JW, Lee BH, Park SG, Kim BC, Lee S, Lee SJ. Factors predicting malignancy in adult intussusception: an experience in university-affiliated hospitals. *Asian J Surg*. 2018;41(1): 92-7.
- Kim KH, Namgung H, Park DG. Adult intus susceptions: preoperative predictive factors for malignant lead point. *Ann Surg Treat Res*. 2014; 86(5): 244–8.