



Sudden Cardiac Deaths Before-After COVID-19 Pandemic and Spike Protein Derivates Uses

Luisetto M^{1*}, Almukhtar N², Khan FA³, Hamid GA⁴, Massimo C⁵, Ansovini R⁶, Tarro G⁷, Cabianca L⁸, Rasool MG⁹, Fiazza C¹⁰, YesviRafa A¹¹, Ahmadabadi NB¹² and Latshev OY¹³

¹General Toxicology, Applied and Industrial Chemistry Branch, IMA Marijnskaya Academy, Italy

²Department of Physiology, College of Medicine, University Babilon, Iraq

³Department of Pharmacology, JN Medical College, India

⁴Department of Hematology Oncology, University of Aden, Yemen

⁵Department of Molecular Biology, Genetic Forensic, Italy

⁶Ansovini Technology, Italy

⁷Department of Oncologic Virology, World Academy of Biomedical Technologies (WABT), France

⁸Medical Laboratory Turin, Citta' della Salute, Italy

⁹Department of Medical and Health Sciences for Woman, Institute of Pharmaceutical Science, Peoples University of Medical and Health Sciences for Women, Pakistan

¹⁰Medical Pharmacology, Independent Researcher, Italy

¹¹University of Nebraska-Lincoln NE, USA

¹²Nano Drug Delivery, USA

¹³President IMA ACADEMY, Russia

Abstract

Aim of this work is to observe some interesting phenomena involving pre- and post-pandemic cardiac sudden deaths epidemiology and after the mass vaccination campaign.

The current level of unexpected extra deaths as reported in scientific literature vs. pre-pandemic years. So, it is need a deeply investigation in order to verify association or not with Spike protein derivates used. Of interest to observe the prevalence/incidence of cardiac sudden deaths after COVID-19 disease before the vaccination seasons.

This review is performed whit a neutral approach and without pre-concept. All data comes from biomedical database and published as article.

Keywords: Cardiac sudden deaths; Myocarditis; Pericarditis; SARS-CoV-2 spike protein derivates; Physiology epidemiology; Pathology; Toxicology

Introduction

Great public debate is involved in the level of extra cardiac sudden deaths as reported in the media in last periods. The same of interest to verify the moves of this phenomena and considering the fact that a severe pandemic (COVID-19-SARS-CoV-2) produced high number of deaths (in acute way) but also as long COVID-19 sequela.

To be considered on the global mortality rate also the effect played by the protocols adopted; Preventive measure (lockdown and other) or therapeutic option (drug used or not used): Interesting literature report the objective analysis in neutral scientific way and the entity of the effect.

The history of this time under evaluation reported various research published or changing the protocols to be used: See the hydroxychloroquine case.

Various effects were due to "Pandemic inside the pandemic". The pathologic effect for the heart and other organ of the virus is due by the SPIKE protein toxicity and by the great immune response of the body to fight this microorganism. The toxicity of the spike protein is clearly showed by scientific literature without any doubt.

The fact that some COVID-19 vaccine use Spike protein derivates (or system to produce inside

OPEN ACCESS

*Correspondence:

Luisetto Mauro, General Toxicology, Applied and Industrial Chemistry Branch, IMA Marijnskaya Academy, Italy, Tel: +39-3402479620; E-mail: mauro65@gmail.com

Received Date: 17 May 2023

Accepted Date: 22 Jun 2023

Published Date: 03 Jul 2023

Citation:

Luisetto M, Almukhtar N, Khan FA, Hamid GA, Massimo C, Ansovini R, et al. Sudden Cardiac Deaths Before-After COVID-19 Pandemic and Spike Protein Derivates Uses. *Ann Clin Case Rep.* 2023; 8: 2433.

ISSN: 2474-1655.

Copyright © 2023 Luisetto M. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1: Cardiovascular complications of COVID-19.

the body this product like mRNA vaccine) to stimulate immune response generate the need to verify its profiles of safety also related cardiac sudden deaths and not only for other risks (central nervous system thrombosis et other).

For this reason, it is useful to observe epidemiology of sudden deaths in year before pandemic and to compare with pandemic season, and verifying also the effect played by long COVID syndrome and if possible, to divide the effect played by the vaccination campaign.

All this factor can be considered as cofactor? And there is a summatory of pathological effect? The rare ADR registered in some m RNA vaccine like myocarditis-pericarditis in young’s and the fact that some authorities restricted use of some product to determinate class age are facts that need to better study these phenomena.

Electronic Devices (WCR) and COVID-19 Vaccine ADR: Myocarditis and Pericarditis –Epidemiology and Physiology of an interesting phenomena.

“After observing this evidence published in scientific article and observing that the median (age) of RARE myocarditis after COVID-19 vaccine seen differ from the classic myocarditis (more frequent in young in vaccinated) and related the profile of the COVID-19 disease (Higher risk of myocarditis >50 year) and the role played by comorbidities it is of interest to match the geographic distribution of myocarditis after COVID-19 vaccine with the use of WCR device. This because these 2 factors (Spike protein natural or artificial) and WCR radiation are recognized able to Influence heart as reported in literature” (Figure 1).

From European Society of Cardiology COVID-19 patients retain elevated risk of death for at least 18 months after infection.

“Compared to uninfected individuals, the risk of death in COVID-19 patients was up to 81-fold higher in the acute phase and 5-fold higher in the post-acute phase. Patients with severe COVID-19 were more likely to develop major CV disease or die than non-severe cases”.

“The results showed that the adjusted incident rate ratios of CV




Figure 2: All-cause excess mortality, COVID-19-associated hospitalisations, and confirmed COVID-19 cases in Massachusetts, USA (A) Shows the cumulative all-cause excess mortality (blue line) plotted against weekly confirmed new COVID-19 cases (light blue line) in Massachusetts, from Feb, 2020, through June, 2022.7 (B) Shows the cumulative number of COVID-19-associated hospitalisations (purple line) plotted against weekly confirmed new COVID-19 cases (light blue line) in Massachusetts, from Feb, 2020, through June, 2022.7 hospitalisation data did not become available until May, 2020.

outcomes in the post-COVID-19 exposure period were significantly higher than those in the pre-exposure period (ratios of incident rate ratios for all CV outcomes were significantly higher than 1) and exhibited a graded increase by severity of the acute phase of the disease. Cardiac arrest (HR=2.45 (2.08-2.89))”.

Figure 2 All-cause excess mortality, COVID-19-associated hospitalizations, and confirmed COVID-19 cases in Massachusetts, USA (A) Shows the cumulative all-cause excess mortality (blue line) plotted against weekly confirmed new COVID-19 cases (light blue line) in Massachusetts, from February, 2020, through June, 2022. 7 (B) Shows the cumulative number of COVID-19-associated hospitalizations (purple line) plotted against weekly confirmed new COVID-19 cases (light blue line) in Massachusetts, from Feb, 2020, through June, 2022; 7 hospitalization data did not become available until May, 2020.

“Myocarditis is a condition caused by acute or chronic inflammation of the cardiac myocytes, resulting in associated myocardial edema and myocardial injury-necrosis. The exact incidence is unknown, but is likely underestimated, with more mild cases going unreported. Diagnosis and appropriate management are paramount in pediatric myocarditis, as it remains a recognized cause of sudden cardiac death SCD in children and athletes. Myocarditis in children is most often caused by a viral or infectious etiology. There are now 2 highly recognized etiologies related to COVID-19 infection and the COVID-19 mRNA vaccine”.




November 27th, 2022

“In general, a causal link between myocarditis and anti-SARS-CoV-2 vaccination is supported by several considerations: A close temporal relation to vaccination; all cases were found dead within 1 week after vaccination, absence of any other significant pre-existing heart disease, especially ischemic heart disease or cardiomyopathy, negative testing for potential myocarditis-causing infectious agents, presence of a peculiar CD4 predominant T-cell infiltrate, suggesting an immune mediated mechanism. The latter criterion is supported by demonstration of a phenotypically identical T-cell infiltrate at the deltoid injection site in 1 of the cases”.

According COVID-19: Pandemic disruption linked to 30,000 excess heart disease deaths, charity reports.

“Between March 2020 and Aug 2022 there were more than 30,000 excess deaths involving coronary heart disease, an average of 230 a week above the expected death rate. Many people have not been able to access care for conditions like as high blood pressure, that could raise the risk of a future heart attack or stroke, the foundation’s report said”.

Excess deaths in 2022 among worst in 50 years “More than 650,000 deaths were registered in the UK in 2022, 9% more than



2019” (Figure 3 and 4).

“SARS-CoV-2 vaccination reduces morbidity and mortality associated with COVID-19 disease; unfortunately, it is associated with serious adverse events AE, including Sudden Unexplained Death (SUD). 13 were recruited, aged between 23 and 72 years; 10 (77%) were men, 12 were Thai; and 1 was Australian. Eight (61%) died after receiving the first dose of vaccine, and 7 (54%) died after receiving ChAdOx1 nCoV-19; There were no significant correlations between SUD and either the number or the type of vaccine. Fever was self-reported in 3 cases. Ten (77%) and 11 (85%) died within 24 h and 3 days of vaccination, respectively. Whole exome sequencing analysis revealed that 5 cases harbored SCN5A variants that had previously been identified in patients with Brugada syndrome BS, giving an SCN5A variant frequency of 38% (5 of 13). This is a significantly higher rate than that observed in Thai SUD cases occurring 8 to 30 days after COVID-19 vaccination during the same period (10% (1 of 10)), in a Thai SUD cohort studied before the COVID-19 pandemic (12% (3 of 25)), and in our in-house exome database (12% (386 of 3231))” (Figure 5).

From MAYO CLINIC:

“Brugada (brew-GAH-dah) syndrome is a rare but potentially

life-threatening heart rhythm condition (arrhythmia) that is sometimes inherited.

People with Brugada syndrome have an increased risk of irregular heart rhythms beginning in the lower chambers of the heart (ventricles)”.

April 28th, 2022

Increased emergency cardiovascular events among under- 40 population in Israel during vaccine rollout and third COVID-19 wave.

“CV adverse outcomes such as blood clotting (coronary artery thrombosis), acute coronary syndrome, cardiac arrest and myocarditis have been identified as consequences of COVID-19 infection. Data from regulatory surveillance and self-reporting systems, including the Vaccine Adverse events Reporting System in the US, the Yellow Card System in the UK and the EudraVigilance system in Europe, associate similar CV side-effects with a number of COVID-19 vaccines currently in use” (Figures 6-8).

From Euroscientist.com:

Science in society » UK recorded more than 10,000 extra non-covid deaths since July: How do we know vaccines are not to blame?





Figure 6: Number of cases of suspected drug- or vaccine-induced myocarditis, perimyocarditis and pericarditis reported to the Regional Pharmacovigilance Centre (RPVC) from 2012 until July 2021. Blue bars and blue lettering indicate the number of myocarditis and perimyocarditis cases, green bars and green lettering indicated the number of pericarditis cases.



Jean-Philippe E, et al. J Am Coll Cardiol. 2022;79(18):1818-1827.

Figure 7: Trends in the incidence of sudden cardiac death.




Figure 8: Weekly mortality in the U.S. reported by the CDC as of October 2022. The red line marks the upper bound of the threshold for expected deaths, above which more deaths than expected have been registered. Weeks of excess mortality are indicated by a red cross.




Figure 9: A wave of unexplained extra deaths from causes other than Covid-19 hit the UK, the first nation to start mass vaccination with novel vaccines.

“Across the UK, more people than usual were dying from non-Covid causes last year—and no one knows why (Figure 9).

Since July 2021, 7 months after the vaccination program started in earnest, over 10,000 more people than usual have died in England and Wales alone from conditions unrelated to coronavirus, according to data from the Office for National Statistics NS. Paul Hunter, professor in medicine at the Univ. of East Anglia, notes that: In general, the excess deaths seem to be primarily CV, and that many of them are happening at home”.

Myocarditis and Pericarditis After Vaccination for COVID-19.

“40 hospitals in Washington, Oregon, Montana, and LA County, California that were part of the Providence health care system and used the same electronic medical record were included. All patients with documented COVID-19 vaccinations administered inside the system or recorded in state registries at any time through May 25th, 2021, were identified. Vaccinated patients who subsequently had Emergency Department ED or inpatient encounters with diagnoses of myocarditis, myopericarditis, or pericarditis were ascertained from EMRs” (Figure 10, 11).

Extra deaths in 2022 close to highest level in 70 years - how much




Figure 10: Incident cases of pericarditis.

are NHS failings to blame?

“More deaths at home, many of the recent excess deaths are occurring at home, as opposed to in hospitals or care homes.

In the 2 weeks to 30th December, 1,804 more people died at home




Figure 11: Weekly counts of cardiac arrest calls (5-week centered moving-average), COVID-19 cases (3-week centered moving-average), and vaccination doses (3-week centered moving-average) for those between 16 and 39 during: A) the study period (Jan 1 2019, to Jun 20, 2021) and B) the third COVID-19 wave and vaccination distribution period (Oct 18, 2020, to Jun 20, 2021) COVID-19 disease 2019.

Many more deaths than usual have occurred at home

Deaths registered between 1 July and 30 December 2022, compared with average from 2015-19



SOURCE: ONS

Figure 12: Many more deaths than usual have occurred at home.




Figure 13: Distribution of events during the day. Bars: average hourly incidence of events computed across different years. Whiskers: standard error. Solid line: moving average.

than usual, almost 40% higher than the 5-year average. In total over the past 6 months 84,337 people have died at home, 21,803 (34.9%) more than the 5-year average from 2015-19. That's a much bigger difference than the change in other settings" (Figure 12).

Materials and Methods

With an observational point of view some relevant literature and Figures 1-23 are reported. An experimental hypothesis is submitted to the researcher in order to provide a global conclusion related the topic of investigation. All literature comes from PubMed or other relevant biomedical database.

Results

From literature

According Cadegiani FA: "Myocarditis and sudden death SD after COVID-19 mRNA vaccination. Myocarditis induced by SARS-CoV-2 mRNA vaccines is an indisputable complication observed particularly in young males as demonstrated by multiple studies in different populations. Two studies reported that COVID-19 mRNA vaccine-induced myocarditis disproportionately affected adolescents (Reporting Odds Ratio (ROR): 22.3; 95% Confidence Interval (CI): 19.2-25.9), 18 to 29-year-olds (ROR: 6.6; 95% CI: 5.9-7.5), and males (ROR: 9.4; 95% CI: 8.3-10.6).

These findings were supported by other great registry study that identified increased myocarditis risk MR following SARS-CoV-2 mRNA vaccination, with the highest risk detected in people aged 18 to 24 years, particularly after 2nd dose, where 8.1-fold increased risk after the BNT162b2 SARS-CoV-2 mRNA vaccine (95% CI: 6.7-9.9) and 30-fold increased risk after the mRNA-1273 SARS-CoV-2 vaccine (95% CI: 21-43) were reported. An Israeli government data set demonstrated 13.6-fold increased myocarditis risk (95% CI: 9.3-19.2) among males aged between 16 to 19 years compared to the expected following historical data, while a nine-fold increased myocarditis risk (95% CI: 4.5-17.8) was recorded when compared to unvaccinated of similar age and sex during the same period. In a 23 million-resident area, the myocarditis risk MR after SARS-CoV-2 mRNA vaccination was increased across all populations, particularly high among males aged 16 to 24 years after the 2nd dose of BNT162b2 and mRNA-1273, where 5.3-fold (95% CI: 3.7-7.7) and 13.8-fold (95% CI: 8.1-23.7) increased myocarditis risk MR was recorded. Independent autopsies

or biopsies of patients who presented post-SARS-CoV-2 mRNA vaccine myocarditis in different geographical regions enabled the conclusion that a primary hyper catecholaminergic state was the key trigger of these events; SARS-CoV-2 mRNA was densely present, and SARS-CoV-2 spike protein was progressively produced in adrenal medulla chromaffin cells, which are responsible for catecholamine production; the dihydroxyphenylalanine decarboxylase enzyme that converts dopamine into noradrenaline was overexpressed in the presence of SARS-CoV-2 mRNA, leading to enhanced noradrenaline activity; catecholamine responses were physiologically higher in young adults and males than in other populations; catecholamine responses and the resting catecholamine production were higher in male athletes than in non-athletes; catecholamine responses to stress and its sensitivity were enhanced in the presence of androgens; and catecholamine expressions in young male athletes were already high at baseline, were higher following vaccination, and were higher than those in non-vaccinated athletes. The epidemiological, autopsy, molecular, and physiological findings unanimously and strongly suggest that a hyper catecholaminergic state is the critical trigger of the rare cases of myocarditis due to components from SARS-CoV-2, potentially increasing sudden deaths SD among elite male athletes. "[1].

"The demographic and clinical characteristics of myocarditis after mRNA vaccination in previous reports are summarized in Table reported. Vaccine-associated myocarditis has been reported predominantly in young males after second vaccination. Myocarditis was mainly diagnosed clinically based on elevated serum troponin and cardiac magnetic resonance imaging, and all patients except ours and the patient reported by Verma et al. Recovered after receiving supportive care [2].

"The incidence of sudden cardiac death ranged from 0.93 to 1.3 per 100,000 person-years, whereas an incidence of 1.62 per 100,000 person-years was reported in this research study [3].

"The study examined 4 large (>2 million inhabitants) European population-based prospective registries collecting emergency medical services-attended (i.e., with attempted resuscitation) OHCA and SCD (OHCA without obvious extracardiac causes) for >5 consecutive years from January 2012 to December 2017 in the Paris region (France), the North Holland region, the Stockholm region (Sweden), and in all of




Figure 14: Several types of CV disease exhibit a time of day incidence in Sudden Cardiac Death (SCD) or events. The left graphic shows the causes and rates for SCD by several different types of CV disease.




Figure 15: Myocarditis Mortality Rate Ratios (MMRRs) by age and Standardised Mortality Ratio (SMR).


Denmark. The average annual incidence of SCD in the 4 registries ranged from 36.8 per 100,000 (95% CI: 23.5-50.1 per 100,000) to 39.7 per 100,000 (95% CI: 32.6-46.8 per 100,000)” [4].

“Cardiovascular CV effects of COVID-19 vaccination

Myocarditis and pericarditis are known complications of mRNA vaccines, especially in young adult and teen males aged 12 to 17 years, 23 with the highest observed incidence within 2 to 7 days after the

second dose at a rate of 3.5-140 per million doses. 27” [5].

“Additional analyses of CDC Vaccine Safety Datalink with data from 9 participating integrated health care organizations revealed an increased risk of myocarditis/pericarditis events among individuals 12 to 39 years of age in the 7-day risk interval after the vaccination with mRNA COVID-19 vaccines compared with unvaccinated individuals or individuals vaccinated with non-mRNA COVID-19



vaccines on the same calendar days (rate ratio of 10.8 [95% CI, 3.2-49.0], adjusted for site, age, sex, race/ethnicity, and calendar date).

The estimated myocarditis/pericarditis chart-confirmed rate was 12.6 cases per million doses with second-dose mRNA vaccine among individuals 12 to 39 years of age. The rates based on International Classification of Diseases; 10th Revision-coded cases were also higher in males than in the females. All chart-confirmed cases with follow-up had resolution of symptoms; and among those who had follow-up ECG/echocardiography and lab. Testing, most had returned to normal or baseline. On this basis, the FDA will add a warning to the product label of both mRNA vaccines regarding the risk of myocarditis” [6].

“In general, treatment of either acute or chronic myocarditis is aimed at reducing congestion and improving cardiac hemodynamics in heart failure HF, as well as providing supportive therapy, with the hope of prolonging survival. Treatment of heart failure follows the same treatment regimen regardless of the underlying cause (ACE inhibitors, beta-adrenergic blockers)” [7].

“We present autopsy findings of a 22-year-old man who developed chest pain 5 days after the first dose of the BNT162b2 mRNA vaccine and died 7 h later. Histological examination of the heart revealed isolated atrial myocarditis, with neutrophil and histiocyte predominance. Immunohistochemical C4d staining revealed scattered single-cell necrosis of myocytes which was not accompanied by inflammatory infiltrates. Extensive contraction band necrosis was observed in the atria and ventricles. There was no evidence of microthrombosis or infection in the heart and other organs. The primary cause of death was determined to be myocarditis, causally-associated with the BNT162b2 vaccine” [2].

“This study focuses on CV manifestation, particularly myocarditis and pericarditis events, after BNT162b2 mRNA COVID-19 vaccine injection in Thai adolescents. This prospective cohort study enrolled students aged 13 to 18 years from 2 schools, who received the second dose of the BNT162b2 mRNA COVID-19 vaccine. Data including demographics, symptoms, vital signs, ECG, echocardiography, and cardiac enzymes were collected at baseline, Day 3, 7, and Day 14 (optional) using case record forms. We enrolled 314 participants; of these, 13 participants were lost to follow-up, leaving 301 participants for analysis. The most common CV signs and symptoms were tachycardia (7.64%), shortness of breath (6.64%), palpitation (4.32%),

chest pain (4.32%), and hypertension (3.99%). One participant could have more than 1 sign and/or symptom. Seven participants (2.33%) exhibited at least 1 elevated cardiac biomarker or positive lab assessments.

CV manifestations were found in 29.24% of patients, ranging from tachycardia or palpitation to myopericarditis. Myopericarditis was confirmed in 1 patient after vaccination. Two patients had suspected pericarditis and 4 patients had suspected subclinical myocarditis. CV manifestation in adolescents after BNT162b2 mRNA COVID-19 vaccination included tachycardia, palpitation, and myopericarditis. The clinical presentation of myopericarditis after vaccination was usually mild and temporary, with all cases fully recovering within 14 days. Adolescents receiving mRNA vaccines should be monitored for CV side effects. Clinical Trial Registration: NCT05288231” [8].

“Cases of myocarditis, diagnosed clinically by laboratory tests and imaging have been described in the context of mRNA-based anti-SARS-CoV-2 vaccination. Autopsy-based description of detailed histological features of vaccine-induced myocarditis is lacking. We describe the autopsy findings and common characteristics of myocarditis in untreated persons who received anti-SARS-CoV-2 vaccination. Standardized autopsies were performed on 25 persons who had died unexpectedly and within 20 days after anti-SARS-CoV-2 vaccination. In 4 patients who received a mRNA vaccination, we identified acute (epi-) myocarditis without detection of another significant disease or health constellation that may have caused an unexpected death. Histology showed patchy interstitial myocardial T-lymphocytic infiltration, predominantly of the CD4 positive subset, associated with mild myocyte damage. Autopsy findings indicated death due to acute arrhythmogenic cardiac failure CF. Myocarditis can be a potentially lethal complication following mRNA-based anti-SARS-CoV-2 vaccination. Our findings may aid in adequately diagnosing unclear cases after vaccination and in establishing a timely diagnosis *in vivo*, providing the framework for adequate monitoring and early treatment of severe clinical cases” [9].

“CV events occur more frequently in the early morning, with morning rates higher for both AMI and sudden cardiac death” [10] (Figure 13).

“A recent analysis of data from Florida (US) found increased risk of cardiac death in the first 4 weeks after mRNA vaccination in people aged 18 to 39 years 20 compared to subsequent weeks” [11].

Exploring the relationship between all-cause and cardiac-related mortality following COVID-19. 2022.

“Cardiac-related deaths CRD following vaccination. In the 28 days following vaccination, a statistically significant increase in cardiac-related deaths was detected for the entire study population (RI=1.07, 95% CI= 1.03-1.12). Stratifying by age group revealed RIs were significantly higher for age groups 25 to 39 (RI=2.16, 95% CI= 1.35-3.47) and 60 or older (RI=1.05, 95% CI=1.01-1.10). The remaining age groups failed to reach statistical significance” [12].

“Myocarditis or pericarditis incidence after mRNA COVID-19 vaccination in the current study (0-35.9 per 100,000 for males and 0-10.9 for females across age groups and vaccine cohorts) was similar to estimates found in a study from eight USA health systems in the Vaccine Safety Datalink [9]. Previous CDC estimates found the highest risk for post-vaccination myocarditis among males aged 16 to 17 years (10.6 per 100,000) during a 7-day risk window after receipt of




Figure 17: A) Circadian variation in the frequency of Sudden Cardiac Death (SCD). The fitted curve is shown in black ($p < 0.01$, $n = 371$). B) Circadian variation in the frequency of non-SCD. The fitted curve is shown in black ($p < 0.01$, $n = 2929$). C) Circadian variation based on fitted curve among SCD and non-SCD. The circadian curve was calculated by (fitted curve value - average fitted curve value) / average fitted curve value.

a second mRNA COVID-19 vaccine dose. Estimates from the current study (22.0 per 100,000 males aged 12 to 17 years) are higher, likely because outcomes were captured using ICD-10-CM codes alone rather than through passive reporting with subsequent verification through medical record review. Among males aged 12 to 17 years, the group with the highest incidence of cardiac complications after receipt of a second mRNA COVID-19 vaccine dose, the risk was 1.8 to 5.6 times as high after SARS-CoV-2 infection than after the vaccination” [13].

“Circadian rhythm and sudden cardiac arrest

In 1987, Muller and his colleagues published a retrospective study of the death certificates of 2203 patients in Massachusetts who died

outside of the hospital in 1983. They found a circadian variation in sudden cardiac death SCD with 2 peaks during 24 h the primary peak occurred from 7 am to 11 am and a secondary peak occurred between 5 pm to 6 pm. These results corresponded with the findings of the MILIS Study Group in 1985 that described the onset of acute myocardial infarction as following a circadian rhythm with a peak from 6 am to noon” [14] (Figure 14).

“This study work compared myocarditis mortality rate in the SARS-CoV-2 vaccinated with that in the general population in Japan. The study was based on the materials and the vital statistics disclosed by the Japanese government.

Number of myocarditis death MD which met the inclusion

criteria were 32 cases. MMRR (95% confidence interval) was 4.03 (0.77 to 13.60) in 20s, 7.80 (2.85 to 18.56) in 30s, respectively. SMR of myocarditis was 1.69 (1.18 to 2.42) for overall vaccinated population, 1.35 (0.84 to 2.55) for those 60 years or older. Estimated adMMRRs and adSMR were about 4 times higher than the MMRRs and SMR. Pooled MOR for myocarditis was 148.49 (89.18 to 247.25).

SARS-CoV-2 vaccination was associated with higher risk of myocarditis death MD, not only in young adults but also in all age groups including the elderly. Considering healthy vaccinee effect, the risk may be 4 times or higher than the apparent risk of myocarditis death. Underreporting should also be considered. Based on this study, risk of myocarditis following SARS-CoV-2 vaccination may be more serious than that reported previously” [15].

Public health Ontario surveillance report

Adverse Events Following Immunization (AEFIs) for COVID-19 in Ontario: Dec 13, 2020 to March 26, 2023

“There have been international reports, including from the US and Israel, of myocarditis and pericarditis following vaccination with COVID-19 mRNA vaccines. Information to date indicates that these events occur more commonly after the second dose, within the week following vaccination (typically within 4-5 days), mainly in adolescents/young adults and more often in males than females” [16] (Figure 15).

“The mean age of the deceased was 62.6 years (age range: 32 to 91 years). Vaxzevria was vaccinated in 9, Comirnaty in 5, Spikevax in 3, and Janssen in 1 person. The time interval between the last vaccination and death ranged from a few minutes (case 17) up to 14 days (case 9). Case 17 involved a person who collapsed in a vaccination center immediately after the vaccination; prompt resuscitation efforts were unsuccessful. A total of 12 deaths occurred at home, 4 deaths at a hospital, 1 death at a vaccination center, and 1 death at work” [17] (Figure 16).

Over half the deaths related to atherosclerosis (13 of 24) on the day of the earthquake occurred between midnight and 6 am, whereas only 3 of 32 (9%) of the deaths during the 7 days before the earthquake occurred at that time of day (P=0.002). The earthquake that struck the Los Angeles area at 4:31 am on January 17th, 1994, was 1 of the strongest earthquakes ever recorded in a major city in North America.

“In a huge analysis of more than 30,000 vaccinated patients who had experienced COVID breakthrough infections (pre-Omicron

variant), scientists found that 6 months later, even the vaccinated incurred a higher risk of death and debilitating long COVID symptoms involving multiple organs (lungs, heart, kidney, brain, others) when compared to controls without evidence of SARS-CoV-2 infection”.

“The post-acute sequelae of SARS-CoV-2 infection- also referred to as Long COVID- have been described, but whether Breakthrough SARS-CoV-2 Infection (BTI) in vaccinated people results in post-acute sequelae is not clear. In this study, we used the US Department of Veterans Affairs national healthcare databases to build a cohort of 33,940 individuals with BTI and several controls of people without evidence of SARS-CoV-2 infection, including contemporary (n=4,983,491), historical (n=5,785,273) and vaccinated (n=2,566,369) controls. At 6 months after infection, we show that, beyond the first 30 days of illness, compared to contemporary controls, people with BTI exhibited a higher risk of death (Hazard Ratio (HR)=1.75, 95% Confidence Interval (CI): 1.59-1.93) and incident post-acute sequelae (HR=1.50, 95% CI: 1.46-1.54), including cardiovascular CV, coagulation and hematologic, GI, kidney, mental health, metabolic, musculoskeletal and neurologic disorders. The results were consistent in comparisons *versus* the historical and vaccinated controls. Compared to people with SARS-CoV-2 infection who were not previously vaccinated (n=113,474), people with BTI exhibited a lower risk of death (HR=0.66, 95% CI: 0.58-0.74) and incident post-acute sequelae (HR=0.85, 95% CI: 0.82-0.89). The findings suggest that vaccination before infection confers only partial protection in the post-acute phase of the disease” [18].

“Approximately 80% of all SCDs occur in the home and around 60% are witnessed” [19] (Figure 17).

Circadian aspects of mortality in hospitalized patients: A retrospective observation from a large cohort

“Circadian Variation in SCD

Circadian patterns have been the subject of considerable interest in CV, including both SCD as well as myocardial infarction MI. Cohen and coworkers analyzed 19 studies on the pattern of incidence of SCD. In their meta-analysis, including 19,390 patients, a morning excess in the incidence of SCD was found. Between 06:00 and 12:00, 30.1% of SCD events occurred; the incidence of SCD was 29% higher in the morning than during the rest of the day (relative risk 1.29; 95%” [20] (Figure 18, 19).

“We find an increase in cardiac death in women after a first dose of non-mRNA vaccines. A positive SARS-CoV-2 test is associated with

TOP FIVE CATEGORIES OF DEATH FOLLOWING VACCINATION					
Diagnosis Category	<3 days	4-7 days	8-28 days	>28 days	Total
Acute Coronary Syndrome/ Myocardial Infarction	49	10	4	0	63
Cerebrovascular Accident	11	2	1	0	14
Sudden Death	7	2	1	0	10
Respiratory Infection	4	0	1	0	5
Unknown	9	2	3	0	14

Source: ITSU AEFI Secretariat presentation to the National AEFI Committee

Figure 18: Top five categories of death following vaccination.




Figure 19: Number of Cardiovascular (CV), trauma-related, and other sudden death events in 1866 young competitive athletes, tabulated by year.

increased cardiac and all-cause mortality among people vaccinated or unvaccinated at time of testing” [21] (Figure 20).

“In the vaccination analysis, the vaccinated and control groups each included a mean of 884,828 persons. Vaccination was most strongly associated with an elevated risk of myocarditis (risk ratio: 3.24; 95% Confidence Interval (CI), 1.55-12.44; risk difference, 2.7 events per 100,000 persons; 95% CI, 1.0-4.6). SARS-CoV-2 infection was associated with a substantially increased risk of myocarditis (risk ratio: 18.28; 95% CI, 3.95-25.12; risk difference, 11.0 events per 100,000 persons; 95% CI, 5.6-15.8) and of additional serious adverse AE events, including pericarditis, arrhythmia, deep-vein thrombosis, pulmonary embolism EP, myocardial infarction, intracranial hemorrhage, and thrombocytopenia. The vaccine was associated with an excess risk of myocarditis (1 to 5 events per 100,000 persons). The risk of this potentially serious adverse event and of many other serious adverse events was substantially increased after SARS-CoV-2 infection” [22].

“There were 19,740,741 doses of mRNA vaccines administered and 297 reports of myocarditis/pericarditis meeting our inclusion criteria. 69.7% occurred following the second dose of COVID-19 mRNA vaccine and 76.8% occurred in males. The median age of individuals with a reported event was 24 years. The highest reporting rate of myocarditis/pericarditis was observed in males aged 18 to 24 years following mRNA-1273 as the second dose; the rate in this age group was 5.1 (95% CI 1.9-15.5) times higher than the rate following BNT162b2 as the second dose. Overall reporting rates were higher when the inter-dose interval was shorter (≤ 30 days) for both vaccine products. Among individuals who received mRNA-1273 for the second dose, rates were higher for those who had a heterologous as opposed to homologous vaccine schedule” [23].

COVID-19 vaccine

Strong association with cardiovascular death, especially

hemorrhagic stroke and venous thrombosis.

“We analyzed 196 death cases reported after inoculation of Pfizer-BioNTech COVID-19 vaccine (COMIRNATY) by June 9 in Japan. Japanese Ministry of Health JMH, Labor, and Welfare virtually denied the causality of all cases without conducting appropriate epidemiological analysis.

Mortality odds ratio was calculated as the indicator of disproportionality in cause of death. We compared odds of cause of death after inoculation and death in Japanese vital statistics in 2019 as control non-vaccinated population. MOR was obtained by using the numbers of death from non-CV system as the reference causes for 2 age groups: Vaccinated medical workers (20 to 74 year of age) and elderly (≥ 65 years) separately.

Of 31 deaths among vaccinated medical workers (both sexes), 26 (84%) died from CV diseases, such as stroke, myocardial infarction, Venous Thrombosis and Pulmonary Embolism (VT/PE) and heart failure, while 22% died in the general population. MOR is 19.4 ($p < 0.0001$). MOR of hemorrhagic stroke (40.7) and VT/PE (114.0) were extremely high.

Of the reported vaccinated elderly death cases, 69% died from CV causes, while 26% died in the general population. MOR is 5.9 ($p < 0.0001$). MOR of hemorrhagic stroke (12.8) and VT/PE (24.9) were also very high.

These suggest that COVID-19 vaccination is closely associated with the risk of death from CV causes, especially hemorrhagic stroke and VT/PE” [24].

But an adverse event occurring after vaccination doesn't prove vaccination caused it (Figure 21).

“During Wave I (1 March 2020 to 6 June 2020), we observed a sharp increase in excess CVD deaths. There was an 8.6% (95%




Figure 20: Relative incidence of cardiac and all-cause death in the risk period after vaccination, sex, age group and vaccine vector.

CI 8.4%; 8.8%) increase in deaths due to Ischemic Heart Disease (IHD) and a 13.2% (95% CI 12.9%; 13.5%) increase in deaths due to hypertensive disease. During Wave II (June 7th, 2020 to October 3rd, 2020), the curves of COVID-19 deaths and excess CVD deaths were both milder. Although there was a large peak of COVID-19 during

Wave III (October 4th, 2020 to June 26th, 2021), the excess CVD deaths were lower compared to those in previous waves” [25].

USA vaccination campaign started on December 2020

“A literature review of COVID-19-vaccine-related deaths has




Figure 21: Comparison of cause of death: Vaccinated and general population.

been carried out according to PRISMA standards to understand if there is a causal relationship between vaccination and death and to highlight the real extent of such events. There have been 55 cases of death after COVID-19 vaccination reported and a causal relationship has been excluded in 17 cases. In the remaining cases, the causal link between the vaccine and the death was not specified (8) or considered possible (15), probable (1), or very probable/demonstrated (14). The causes of deaths among these cases were: Vaccine-induced immune thrombotic thrombocytopenia (32), myocarditis (3), ADEM (1), myocardial infarction (1), and rhabdomyolysis [26] (Figure 22, 23).

“In this study work in a nationwide mass vaccination setting, the BNT162b2 vaccine was not associated with an elevated risk of most of the adverse events examined. The vaccine was associated with an excess risk of myocarditis (1 to 5 events per 100,000 persons). The risk of this potentially serious adverse event and of many other

serious adverse events was substantially increased after SARS-CoV-2 infection” [27].

“62 studies, including 218 cases, participated in the current systematic review. The median age was 29.2 years; 92.2% were male and 7.8% were female. 72.4% of patients received the Pfizer-BioNTech (BNT162b2) vaccine, 23.8% of patients received the Moderna COVID-19 Vaccine (mRNA-1273), and the rest of the 3.5% received other types of COVID-19 vaccine. Most myocarditis cases (82.1%) occurred after the second vaccine dose, after a median time interval of 3.5 days. The most frequently reported symptoms were chest pain, myalgia/body aches and fever. The Troponin levels were consistently elevated in 98.6% of patients. The admission ECG was abnormal in 88.5% of cases, and the left LVEF was lower than 50% in 21.5% of cases. Most patients (92.6%) resolved symptoms and recovered and only 3 patients died” [28].

“The Israeli Ministry of Health reported 62 cases of myocarditis in patients vaccinated for COVID-19 out of 5 million vaccinated individuals. Great number cases occurred after the second dose of mRNA vaccines, with only 6 cases diagnosed after the first dose. The prevalence was higher in men under 30 years of age, increasing from 1/100 000 for the general population, to 1/20 000 for the 16 to 30 years old group. Two of the 62 patients died. The US Department of Defense reported 14 military personnel diagnosed with myocarditis following COVID vaccination, 13 of them after their second dose of COVID-19 mRNA vaccines” [29].

Experimental project hypothesis

1. Because the toxicity of SARS-CoV-2 spike protein and its derivatives is known it is of interest to verify. The incidence and




Figure 22: From pandemic disrupted historical mortality patterns, caused largest jump in deaths in 100 Years.




Figure 23: Average daily deaths by month in the United States: 2010-2021.

prevalence of sudden cardiac deaths in a determinate geographic population before COVID-19 pandemic

2. During pandemic and after COVID disease with vaccination
3. During pandemic after COVID-19 disease and with vaccination
4. After pandemic with only vaccination (no disease)

This data makes possible to verify the contribution of every single move.

Discussion

The profile of CV complication of COVID-19 disease is clearly reported by literature. Related reference reported it is possible to verify this temporal presentation of effects: The same the infected patient (with severe disease) shows increased level of cardiovascular complication vs. the non-infected. (ESC 2023) This data was higher vs. pre-pandemic periods [18]. In Figure 20 the monthly death 2020-21 was always superior than in 2018-19 years even if after starting the mass vaccinations. The same the monthly death in years from 2010 to 2019 was lower than 2020-21. (Figure 23). Great part of sudden deaths are of cardiovascular origin. Extra deaths are reported from January 2021. A part from COVID-19 disease deaths are registered extra deaths. (February-July 2022 UK- Figure 3). Pericarditis increased from May 2021 (USA) Figure 10. Cardiac arrest call reported peaks on 2021 March and July (Figure 11).

Of interest the circadian profile of CSD and the days of presentation after vaccination/when happened and registered).

Great part of deaths after COVID-19 vaccination happened in

less than 3 days (Figure 18).

Of interest also to observe the pathological mechanism of action of SARS-CoV-2 spike protein through the link with ACE2 receptor in pulmonary and other tissue. The same to observe the RARE ADR of some mRNA COVID-19 vaccine like pericarditis and myocarditis in young males. Of interest to observe the role played by ACE Inhibitors in myocarditis therapy. The incidence of CV Deaths or unexpected deaths before and after the COVID-19 vaccination campaign is an interesting phenomenon.

Trends from 2012 to 2017 are reported in references (not great increased), Figure 7. Related the data published it is possible to consider that the similar population pre and post-pandemic studied have an equal level of basic cardiovascular pathology, so the additional mortality can not be considered due only by this phenomena (Figure 23 related daily deaths).

To be considered also that various references report that after vaccination some events are registered in healthy patients. (Are there relationship?) [29], military often are healthy. Some producers of COVID-19 vaccine introduced in COVID vaccine-technical sheet information about risk of rare myocarditis in subpopulations. Relevant the study related the SUD in Brugada syndrome associated a COVID-19 vaccination (genetic factors involved) Figure 5. The vaccination after the COVID disease can be considered like and additional dose of Spike protein ([5] for the cumulative effect).

The incidence of myocarditis after some COVID-19 vaccination was higher in some subclasses (young then adults).

Event like pericarditis and myocarditis increase after the second dose rates were higher when the inter-dose interval was shorter (i.e., ≤ 30 days). Of interest to observe that mortality increase in statistical

way every year due by the difference between new born and new elderly (so increase elderly and related deaths). The average data of increase is greater than of the 2015-19. The difference between 2022 and 2021 can be used a measure of the vaccine influence. In the 2022 it was observed a +12% vs. 2021. The fact that in 2020 was reported more deaths can be related to various factors (Pandemic, kind of preventive measures adopted and kind of therapy chosen). Literature published provided evidence of this, if mortality in 2020 was only due to SARS-CoV-2 virus and no other co-factor (also by some kind of preventive measure, drugs used or not, protocol adopted in emergency) in the 2021-2022 it had to be observed to a reduction: Instead, the cause of death for all causes was increased.

Conclusion

Even Considering that the prophylaxis measure like vaccination is useful in various virus disease observing scientific database some questions are of interest. For the meaning of this work, it is necessary read the data and figure related CSD or deaths considering pre-pandemic years, pandemic season and also the vaccine campaign timing (first doses and second or other doses). Why monthly mortality rate was higher in 2021 than in 2013 in USA? What is the cumulative effect played on Sudden cardiac extra deaths and unexpected by COVID disease or long COVID added to vaccination? And due only by vaccination? It is possible to choose 2013 data as baseline. This information is useful to better understand these global phenomena. Why great amount of unexpected extra death was observed at home? The figure reported in this work added to the time presentation of the effect can give information about these phenomena.

The fact that was registered extra deaths in last periods a part the classic parameter (2012-2017) can be due by the effect of the pandemic (or to long COVID) but it must to be more deeply investigated also the effect played by the spike protein derivatives (wide used) to fight this virus.

References

- Cadegiani FA. Catecholamines are the key trigger of COVID-19 mRNA vaccine-induced myocarditis: A compelling hypothesis supported by epidemiological, anatomopathological, molecular, and physiological findings. *Cureus*. 2022;14(8):e27883.
- Choi S, Lee SH, Seo JW, Kim MJ, Jeon YH, Park JH, et al. Myocarditis-induced sudden death after BNT162b2 mRNA COVID-19 vaccination in Korea: Case report focusing on histopathological findings. *J Korean Med Sci*. 2021;36(40):e286.
- Vaartjes I, Hendrix A, Hertogh EM, Grobbee DE, Doevendans PA, Mosterd A, et al. Sudden death in persons younger than 40 years of age: Incidence and causes. *Eur J Cardiovasc Prev Rehabil*. 2009;16(5):592-6.
- Empana JP, Lerner I, Valentin E, Folke F, Böttiger B, Gislason G, et al. Incidence of sudden cardiac death in the European Union. *J Am Coll Cardiol*. 2022;79(18):1818-27.
- Akhtar Z, Trent M, Moa A, Tan TC, Fröbert O, MacIntyre CR. The impact of COVID-19 and COVID vaccination on cardiovascular outcomes. *Eur Heart J Suppl*. 2023;25(Suppl A):A42-9.
- Bozkurt B, Kamat I, Hotez PJ. Myocarditis with COVID-19 mRNA vaccines. *Circulation*. 2021;144:471-84.
- Tang WHW. Myocarditis Medication. *Medscape*. 2021.
- Mansangan S, Charunwatthana P, Piyaphanee W, Dechkhajorn W, Poolcharoen A, Mansangan C. Cardiovascular manifestation of the BNT162b2 mRNA COVID-19 vaccine in adolescents. *Trop Med Infect Dis*. 2022;7(8):196.
- Schwab C, Domke LM, Hartmann L, Stenzinger A, Longereich T, Schirmacher P. Autopsy-based histopathological characterization of myocarditis after anti-SARS-CoV-2-vaccination. *Clin Res Cardiol*. 2023;112(3):431-40.
- Peñaloza-Martínez E, Moreno G, Aroca-Crevillén A, Huertas S, Vicent L, Rosillo N, et al. Circadian rhythms in thrombosis and atherothrombotic events. *Front Biosci (Landmark Ed)*. 2022;27(2):51.
- Nafilyan V, Bermingham CR, Ward IL, Morgan J, Zaccardi F, Khunti K, et al. Risk of death following COVID-19 vaccination or positive SARS-CoV-2 test in young people in England. *Nat Commun*. 2023;14(1):1541.
- Exploring the relationship between all-cause and cardiac-related mortality following COVID-19. 2022.
- Block JP, Boehmer TK, Forrest CB, Carton TW, Lee GM, Ajani UA, et al. Cardiac complications after SARS-CoV-2 infection and mRNA COVID-19 vaccination - PCORnet, United States, January 2021-January 2022. *MMWR Morb Mortal Wkly Rep*. 2022;71(14):517-23.
- Ramireddy A, Chugh SS. Do peak times exist for sudden cardiac arrest? *Trends Cardiovasc Med*. 2021;31(3):172-6.
- Watanabe S, Hama R. SARS-CoV-2 vaccine and increased myocarditis mortality risk: A population based comparative study in Japan. *MedRxiv*. 2022.
- Adverse Events Following Immunization (AEFIs) for COVID-19 in Ontario: December 13, 2020 to March 26, 2023. *Public Health Ontario Surveillance Report*. 2023.
- Schneider J, Sottmann L, Greinacher A, Hagen M, Kasper HU, Kuhnen C, et al. Postmortem investigation of fatalities following vaccination with COVID-19 vaccines. *Int J Legal Med*. 2021;135(6):2335-45.
- Al-Aly Z, Bowe B, Xie Y. Long COVID after breakthrough SARS-CoV-2 infection. *Nat Med*. 2022;28(7):1461-7.
- Adabag AS, Luepker RV, Roger VJ, Gersh BJ. Sudden cardiac death: Epidemiology and risk factors. *Nat Rev Cardiol*. 2010;7(4):216-25.
- Mahmoud KD, de Smet BJGL, Zijlstra F, Rihal CS, Holmes DR Jr. Sudden cardiac death: Epidemiology, circadian variation, and triggers. *Curr Probl Cardiol*. 2011;36(2):56-80.
- Nafilyan V, Bermingham CR, Ward IL, Morgan J, Zaccardi F, Khunti K, et al. Risk of death following COVID-19 vaccination or positive SARS-CoV-2 test in young people in England. *Nat Commun*. 2023;14(1):1541.
- Branda N, Dagan N, Ben-Shlomo Y, Kepten E, Waxman J, Ohana R, et al. Safety of the BNT162b2 mRNA COVID-19 vaccine in a nationwide setting. *N Engl J Med*. 2021;385:1078-90.
- Buchan SA, Seo CY, Johnson C, Alley S, Kwong JC, Nasreen S, et al. Epidemiology of myocarditis and pericarditis following mRNA vaccination by vaccine product, schedule, and interdose interval among adolescents and adults in Ontario, Canada. *JAMA Netw Open*. 2022;5(6):e2218505.
- COVID-19 vaccine: Strong association with cardiovascular death especially hemorrhagic stroke and venous thrombosis. *Med Check Editorial Team*.
- Han L, Zhao H, Li S, Gu S, Deng X, Yang L, et al. Excess cardiovascular mortality across multiple COVID-19 waves in the United States from March 2020 to March 2022. *Nat Cardio Vasc Res*. 2023;2:322-33
- Maiese A, Baronti A, Manetti AC, Paolo MD, Turillazzi E, Frati P, et al. Death after the administration of COVID-19 vaccines approved by EMA: Has a causal relationship been demonstrated? *Vaccines (Basel)*. 2022;10(2):308.
- Barda N, Dagan N, Ben-Shlomo Y, Kepten E, Waxman J, Ohana R, et al. Safety of the BNT162b2 mRNA COVID-19 vaccine in a nationwide setting. *N Engl J Med*. 2021;385:1078-90.
- Ahmed SK, Mohamed MG, Essa RA, Rashad EAA, Ibrahim PK, Khdir AA, et al. Global reports of myocarditis following COVID-19 vaccination:

A systematic review and meta-analysis. *Diabetes Metab Syndr.* 2022;16(6):102513.

29. Albert E, Aurigemma G, Saucedo J, Gerson DS. Myocarditis following COVID-19 vaccination. *Radiol Case Rep.* 2021;16(8):2142-5.